## Information to support decision making

#### Information for patients, families and the public

The Faculty of Intensive Care Medicine have produced <u>a guide to best practice and</u> <u>decision-making in critical care at the end of life</u>, for patients, relatives and the public on care at the end of life.

Key questions to consider are:

- How will critical care treatments help the person in the short and long term?
- Could critical care treatments offer a quality of life that is acceptable to the person?
- Could critical care treatments help achieve a patient's goals for a good life?
- Are there non-critical-care treatments that may help the person and be more comfortable for them?

#### ICNARC data on outcomes for COVID-19

ICNARC (the Intensive Care National Audit and Research Centre) has published a <u>COVID-19 report</u> on the first reported patients critically ill with COVID-19. See the <u>ICNARC website</u> for the latest version.

### ICNARC data on outcomes for pneumonia

ICNARC has mortality data at discharge from the critical care unit and at ultimate discharge from acute hospital (excluding readmissions of the same patient within the same acute hospital stay) for different age groups.

Note: the data below is for all-cause pneumonia, not COVID-19.

#### **Overall mortality data**

| Age (years) | Critical care unit mortality | Acute hospital mortality |  |
|-------------|------------------------------|--------------------------|--|
| 16-49       | 1,858/16,700 (11.1%)         | 2,244/15,305 (14.7%)     |  |
| 50-59       | 2,527/13,075 (19.3%)         | 3,090/11,913 (25.9%)     |  |
| 60-69       | 4,516/17,835 (25.3%)         | 5,471/16,257 (33.7%)     |  |
| 70-79       | 6,052/19,371 (31.2%)         | 7,447/17,567 (42.4%)     |  |
| 80+         | 3,736/10,183 (36.7%)         | 4,858/9,466 (51.3%)      |  |
| All ages    | 18,689/77,164 (24.2%)        | 23,110/70,508 (32.8%)    |  |
|             |                              | © ICNARC 2020            |  |

# Data for patients with cardiovascular co-morbidity

| Age (years) | Critical care unit<br>mortality - with<br>cardiovascular<br>comorbidity  | Critical care unit<br>mortality - without<br>cardiovascular<br>comorbidity | Acute hospital<br>mortality - with<br>cardiovascular<br>comorbidity | Acute hospital<br>mortality - without<br>cardiovascular<br>comorbidity |
|-------------|--|--|---|--|
| 16-49       | 25/135 (18.5%)   | 1,829/16,503<br>(11.1%)  | 35/128 (27.3%)  | 2,204/15,118<br>(14.6%)  |
| 50-59       | 58/170 (34.1%)   | 2,459/12,871<br>(19.1%)  | 63/157 (40.1%)  | 3,017/11,722<br>(25.7%)  |
| 60-69       | 112/335 (33.4%)  | 4,395/17,463<br>(25.2%)  | 133/307 (43.3%)   | 5,329/15,914<br>(33.5%)  |
| 70-79       | 195/486 (40.1%)  | 5,842/18,844<br>(31.0%)  | 241/430 (56.0%)   | 7,189/17,098<br>(42.0%)  |
| 80+         | 148/338 (43.8%)  | 3,575/9,822<br>(36.4%)   | 195/312 (62.5%)   | 4,648/9,131<br>(50.9%)   |
| All ages    | 538/1,464<br>(36.7%)   | 18,100/75,503<br>(24.0%)   | 667/1,334 (50.0%)   | 22,387/68,983<br>(32.5%)   |
| Definition: | Severe comorbidities must have been present in the 6 months prior to admission to the critical care unit and documented in the patient record either at or prior to admission to the critical care unit. |  |   |  |
|             | Cardiovascular comorbidity is defined as fatigue, claudication, dyspnoea or angina at rest due to myocardial or peripheral vascular disease (New York Heart Association functional class IV).            |  |   |  |
|             | © ICNARC 2020  |  |   |  |

## Data for patients with respiratory co-morbidity

| Age (years) | Critical care unit mortality - with respiratory comorbidity  | Critical care unit<br>mortality - without<br>respiratory<br>comorbidity | Acute hospital<br>mortality - with<br>respiratory<br>comorbidity | Acute hospital<br>mortality - without<br>respiratory<br>comorbidity |  |
|-------------|--|---|--|---|--|
| 16-49       | 121/796<br>(15.2%)   | 1,733/15,842<br>(10.9%)   | 148/745 (19.9%)  | 2,091/14,501<br>(14.4%)   |  |
| 50-59       | 187/668<br>(28.0%)   | 2,330/12,373<br>(18.8%)   | 219/635 (34.5%)  | 2,861/11,244<br>(25.4%)   |  |
| 60-69       | 382/1,171<br>(32.6%)   | 4,125/16,627<br>(24.8%)   | 464/1,109 (41.8%)  | 4,998/15,112<br>(33.1%)   |  |
| 70-79       | 464/1,158<br>(40.1%)   | 5,573/18,172<br>(30.7%)   | 554/1,087 (51.0%)  | 6,876/16,441<br>(41.8%)   |  |
| 80+         | 173/415<br>(41.7%)   | 3,550/9,745<br>(36.4%)  | 221/391 (56.5%)  | 4,622/9,052<br>(51.1%)  |  |
| All ages    | 1,327/4,208<br>(31.5%)   | 17,311/72,759<br>(23.8%)  | 1,606/3,967<br>(40.5%)   | 21,448/66,350<br>(32.3%)  |  |
| Definition: | Severe comorbidities must have been present in the 6 months prior to admission to the critical care unit and documented in the patient record either at or prior to admission to the critical care unit. |   |  |   |  |
|             | Respiratory comorbidity is defined as permanent shortness of breath with light activity due to pulmonary disease or use of home ventilation (excluding CPAP for sleep apnoea).                           |   |  |   |  |
|             |  |   |  | © ICNARC 2020   |  |

| © NICE 2020. All rights reserved. Subject to Notice of rights. |
|--|
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |
|  |